
Athletic Physical Examination Form

Name:			E	Birth Date:		Gender:	M	F		
Grade:		School:	Sport:							
				Home Phone:						
				Daytime Phone:						
				Daytime Phone:						
Additional Emerge										
Name:		Rela	tion:		Dayl	time Phor	ie:			
Medical Hi	story									
			Yes	No				Yes	No	
Any significant pa	st injuries	-			Hospitalizati	ions or su	rgeries			
Allergies, asthma,		ng			Seizures					
Contact lenses or glasses					Head injuries or concussions					
Currently on med	ication/me	dications			Bone or joint injuries					
Chronic illness					Current on all vaccinations					
Allergies					Other:					
Physical E	xam									
	Result	Comments				Result	Comments			
Ears				Neurological						
Nose				Genito-urinary						
Throat				Gastrointestinal						
Eyes				Spinal						
Skin				Mental Health						
Dental/Mouth					diovascular			(1992)		
Lungs				Mus	culoskeletal					
Final Diagnosis:							Yes	No	_	
Physician/PNP Nar	ne:									
Physician/PNP Sigr	nature:						Date:			



Bridgeport Catholic Academy

HEADS UP CONCUSSION ACTION PLAN



IF YOU SUSPECT THAT AN ATHLETE HAS A CONCUSSION, YOU SHOULD TAKE THE FOLLOWING STEPS:

- 1. Remove the athlete from play.
- Ensure that the athlete is evaluated by a health care professional experienced in evaluating for concussion. Do not try to judge the seriousness of the injury yourself.
- Inform the athlete's parents or guardians about the possible concussion and give them the fact sheet on concussion.
- 4. Keep the athlete out of play the day of the injury. An athlete should only return to play with permission from a health care professional, who is experienced in evaluating for concussion.
- "IT'S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON."

Athletes who experience one or more of

CONCUSSION SIGNS AND


Athletes who experience one or more of the signs and symptoms listed below after a bump, blow, or jolt to the head or body may have a concussion.

SYMPTOMS REPORTED BY ATHLETE

- · Headache or "pressure" in head
- · Nausea or vomiting
- · Balance problems or dizziness
- Double or blurry vision
- · Sensitivity to light
- · Sensitivity to noise
- · Feeling sluggish, hazy, foggy, or groggy
- · Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

SIGNS OBSERVED BY COACHING STAFF

- Appears dazed or stunned
- · Is confused about assignment or position
- · Forgets an instruction
- · Is unsure of game, score, or opponent
- Moves clumsily
- · Answers questions slowly
- Loses consciousness (even briefly)
- · Shows mood, behavior, or personality changes
- · Can't recall events prior to hit or fall



JOIN THE CONVERSATION AT www.facebook.com/CDCHeadsUp

HEADS UP

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the

Online Concussion Training for Coaches and Athletic Directors

Online training in sports concussion for youth sports coaches https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2904626/

Free Concussion Training - National Alliance for Youth Sports

https://www.nays.org/resources/more/free-concussion-training/

Concussions in Youth Sports Training Course - NYSCA Online

https://www.nays.org/nyscaonline/preview/concussiontraining.cfm

Concussion Information Sheet

What can happen if my child keeps on playing with a concussion or returns too soon?

Athletes with the signs and symptoms of concussion should be removed from play immediately. Continuing to play with the signs and symptoms of a concussion leaves the young athlete especially vulnerable to greater injury. There is an increased risk of significant damage from a concussion for a period of time after that concussion occurs, particularly if the athlete suffers another concussion before completely recovering from the first one. This can lead to prolonged recovery, or even to severe brain swelling (second impact syndrome) with devastating and even fatal consequences. It is well known that adolescent or teenage athletes will often fail to report symptoms of injuries. Concussions are no different. As a result, education of administrators, coaches, parents and students is the key to student-athlete's safety.

If you think your child has suffered a concussion

Any athlete even suspected of suffering a concussion should be removed from the game or practice immediately. No athlete may return to activity after an apparent head injury or concussion, regardless of how mild it seems or how quickly symptoms clear, without medical clearance. Close observation of the athlete should continue for several hours. The Return-to-Play Policy of the IESA and IHSA requires athletes to provide their school with written clearance from either a physician licensed to practice medicine in all its branches or a certified athletic trainer working in conjunction with a physician licensed to practice medicine in all its branches prior to returning to play or practice following a concussion or after being removed from an interscholastic contest due to a possible head injury or concussion and not cleared to return to that same contest. In accordance with state law, all schools are required to follow this policy.

You should also inform your child's coach if you think that your child may have a concussion. Remember it's better to miss one game than miss the whole season. And when in doubt, the athlete sits out.

For current and up-to-date information on concussions you can go to: http://www.cdc.gov/ConcussionInYouthSports/

Student/Parent Consent and Acknowledgements

By signing this form, we acknowledge we have been provided information regarding concussions.

Student

Student Name (Print):	Grade:
Student Signature:	Date:
Parent or Legal Guardian	
Name (Print):	<u> </u>
Signature:	Date:
Relationship to Student:	

Each year IESA member schools are required to keep a signed Acknowledgement and Consent form and a current Pre-participation Physical Examination on file for all student athletes.

Concussion Information Sheet

A concussion is a brain injury and all brain injuries are serious. They are caused by a bump, blow, or jolt to the head, or by a blow to another part of the body with the force transmitted to the head. They can range from mild to severe and can disrupt the way the brain normally works. Even though most concussions are mild, all concussions are potentially serious and may result in complications including prolonged brain damage and death if not recognized and managed properly. In other words, even a "ding" or a bump on the head can be serious. You can't see a concussion and most sports concussions occur without loss of consciousness. Signs and symptoms of concussion may show up right after the injury or can take hours or days to fully appear. If your child reports any symptoms of concussion, or if you notice the symptoms or signs of concussion yourself, seek medical attention right away.

Symptoms may include one or more of the following:

- Headaches
- "Pressure in head"
- Nausea or vomiting
- Neck pain
- Balance problems or dizziness
- Blurred, double, or fuzzy vision
- Sensitivity to light or noise
- Feeling sluggish or slowed down
- Feeling foggy or groggy
- Drowsiness
- Change in sleep patterns

- Amnesia
- "Don't feel right"
- Fatigue or low energy
- Sadness
- Nervousness or anxiety
- Irritability
- More emotional.
- Confusion
- Concentration or memory problems (forgetting game plays)
- Repeating the same question/comment

Signs observed by teammates, parents and coaches include:

- Appears dazed
- Vacant facial expression
- Confused about assignment
- Forgets plays
- Is unsure of game, score, or opponent
- Moves clumsily or displays in coordination
- Answers questions slowly
- Slurred speech
- Shows behavior or personality changes
- Can't recall events prior to hit
- Can't recall events after hit
- Seizures or convulsions
- Any change in typical behavior or personality
- Loses consciousness